

Notes on anomalies, baryons, and Seiberg duality

Steven Corley

Department of Physics
Brown University
Providence, RI 02912
scorley@het.brown.edu

Abstract

We consider an $\mathcal{N} = 1$ $SU(N_c)$ SUSY gauge theory with $N_f \geq N_c$ matter multiplets transforming in the fundamental and antifundamental representations of the gauge group. Using the Konishi anomaly and a non-anomalous conservation law, we derive a system of partial differential equations that determine the low energy effective superpotential as a function of the mesonic and baryonic vacuum expectation values. We apply the formalism to the cases of $N_f = N_c$ and $N_f = N_c + 1$ where the equations are easily integrated and recover the known results. We further apply the formalism to derive a system of partial differential equations to determine the low energy effective superpotential for the Seiberg dual theories. Finally we briefly discuss the associated matrix models via the Dijkgraaf-Vafa conjecture.

1 Introduction

Quantum field theories with $\mathcal{N} = 1$ supersymmetry were studied in great detail in the mid 90's, for reviews see eg. [1, 2]. One of the problems addressed was that of the dynamically generated superpotential. Seiberg [3] showed that one can go a long way in determining the exact form of the superpotential for various theories using symmetries and holomorphy.

Recently a new technique for evaluating low energy effective superpotentials has been developed by Dijkgraaf and Vafa [4, 5, 6]. They conjectured that the exact low energy effective superpotential for an $\mathcal{N} = 1$ SUSY gauge theory with an adjoint matter field could be constructed from an associated matrix model. The action of the matrix model consists of the superpotential appearing in the definition of the gauge theory and the variables of the matrix model are the superfields themselves. The conjecture then relates the planar diagrammatic contribution to the free energy of the matrix model to the exact low energy effective superpotential of the corresponding gauge theory.

Three different proofs of the conjecture have been given, [7, 8, 9]. [7] treated the theory as an $\mathcal{N} = 2$ theory broken to $\mathcal{N} = 1$ by the presence of a superpotential and then analyzed the gauge theory using the Seiberg-Witten formalism [10, 11]. The low energy effective superpotential was constructed and shown to be the same as arising from the associated matrix model via the Dijkgraaf-Vafa conjecture. In [8] the gauge theory was analyzed perturbatively with the result that the diagrams relevant to the low energy effective superpotential were indeed planar and corresponded directly to planar graphs of the associated matrix model. In [9] a generalized Konishi anomaly equation was constructed which takes the exact form of a matrix model loop equation. The solution of the anomaly equations was shown to correspond to the solution of the associated matrix model loop equations in exactly the way conjectured by Dijkgraaf and Vafa.

The purpose of this paper is to analyze the $\mathcal{N} = 1$ SUSY gauge theory with fundamental matter only along the lines of the Konishi anomaly analysis of [9]. This particular topic has been studied in [12, 13, 14, 15, 16, 17]. Specifically we shall be interested in theories with the number of flavors satisfying $N_f \geq N_c$ and vanishing superpotential. For such theories the low energy degrees of freedom consists of mesons and baryons. For $N_f = N_c$ and $N_f = N_c + 1$ the exact results are known, [18]. In the former case there is no dynamically generated superpotential but the classical constraint equation on the meson and baryon vacuum expectation values is modified. In the latter case there is a dynamically generated superpotential which serves to enforce the classical constraint equations on the meson and baryon vevs.

The remaining cases $N_f \geq N_c + 2$ are less well understood. For $3N_c < N_f$ the theory is infrared free while for $3/2N_c \leq N_f \leq 3N_c$ the theory is believed to be a conformal field theory. In the range $N_c + 1 < N_f < 3/2N_c$ the theory can no longer be conformal as such an assumption would violate unitarity. Instead Seiberg has conjectured [19] that the theory has a dual description in terms of an $SU(N_f - N_c)$ gauge theory with N_f fundamental matter multiplets along with a set of scalar multiplets. This conjecture has passed various consistency checks including showing that both theories have the same

global symmetries, have matching quantum numbers for the gauge invariant operators, and have matching anomalies.

By deriving the Konishi anomaly equation, along with a non-anomalous conservation law, we construct a set of partial differential equations for the low energy effective superpotential W_{eff} as a function of the meson and baryon operators for $N_f \geq N_c$. The elementary superpotential of the theory consists of terms giving vevs to the mesons and baryons of the theory enforced by introducing Lagrange multiplier fields. The Konishi anomaly and conservation equations are enough to solve for the Lagrange multipliers in terms of the meson and baryon operators. These solutions are then used to construct a set of partial differential equations for W_{eff} determining it up to a function of the glueball superfield.

In sections 2 and 3 we present the derivations and results for the $N_f = N_c$ and $N_f = N_c + 1$ cases respectively. In these cases we can solve the equations completely and are able to recover the results of [18]. In sections 4 and 5 we derive the Konishi anomaly and conservation equations for the theory and its Seiberg dual respectively for $N_f \geq N_c + 2$. In this case the equations become considerably more complicated and we are unable to solve them. Nevertheless the set of equations could in principle be used to provide a proof of Seiberg's conjectured duality. In section 5 we consider the associated matrix model and derive its loop equations.

2 $N_f = N_c$

The situation with $N_f = N_c = N$ is somewhat simpler, so we begin with it. As our superpotential we shall simply take

$$W = m_{\tilde{f}}^f (\hat{M}_{\tilde{f}}^f - M_{\tilde{f}}^f) + b(\hat{B} - B) + \tilde{b}(\hat{\tilde{B}} - \tilde{B}) \quad (1)$$

where we have defined the meson and baryon operators

$$\hat{M}_{\tilde{f}}^f := \tilde{Q}_{\tilde{f}}^a Q_a^f \quad (2)$$

$$\hat{B} := \det Q = \frac{1}{N!} \epsilon_{f_1 \dots f_N} \epsilon^{a_1 \dots a_N} Q_{a_1}^{f_1} \dots Q_{a_N}^{f_N} \quad (3)$$

$$\hat{\tilde{B}} := \det \tilde{Q} = \frac{1}{N!} \epsilon^{\tilde{f}_1 \dots \tilde{f}_N} \epsilon_{a_1 \dots a_N} \tilde{Q}_{\tilde{f}_1}^{a_1} \dots \tilde{Q}_{\tilde{f}_N}^{a_N}. \quad (4)$$

The fields $m_{\tilde{f}}^f$, b , and $\tilde{b}$ are Lagrange multipliers that enforce the conditions that the vacuum expectation values of the meson and baryon fields are given respectively by $M_{\tilde{f}}^f$, B and $\tilde{B}$. In the context of matrix models and the Dijkgraaf-Vafa conjecture, this kind of superpotential was introduced in [20] for the mesons and generalized to include baryons in [21]. This theory has an $SU(N_f)_Q \times SU(N_f)_{\tilde{Q}} \times U(1)_R \times \prod_{i=1}^{N_f} U(1)_{i,Q} \times \prod_{i=1}^{N_f} U(1)_{i,\tilde{Q}}$ symmetry¹ with the transformation properties of the fields given in the table below.

¹Note that the $U(1)$ flavor symmetries are not all independent because of the $SU(N_f)$ symmetries.

	$SU(N_f)_Q$	$SU(N_f)_{\tilde{Q}}$	$U(1)_{i,Q}$	$U(1)_{i,\tilde{Q}}$	$U(1)_R$
$Q^f :$	N_f	1	δ_i^f	0	$(N_f - N_c)/N_f$
$\tilde{Q}_{\tilde{f}} :$	1	$\bar{N}_f$	0	$\delta_{\tilde{f}}^i$	$(N_f - N_c)/N_f$

If there is no superpotential then the flavor symmetries would classically give rise to the conservation equations $\bar{D}^2(Q_f^\dagger e^V Q^{f'}) = 0$ and $D^2(\tilde{Q}_{\tilde{f}}^{\dagger\dagger} e^V \tilde{Q}_{\tilde{f}}) = 0$. The $U(1)$ symmetries are however anomalous and are modified at one-loop [22, 23] to

$$\bar{D}^2(Q_{f'}^\dagger e^V Q^f) = S \delta_{f'}^f \quad (5)$$

$$\bar{D}^2(\tilde{Q}_{\tilde{f}} e^V \tilde{Q}^{\tilde{f}'\dagger}) = S \delta_{\tilde{f}}^{\tilde{f}'} \quad (6)$$

respectively, where the glueball superfield S is defined as

$$S := \frac{1}{32\pi^2} \text{Tr} W^\alpha W_\alpha. \quad (7)$$

Reinstating the superpotential (1), the anomaly equations pick up some extra terms from the equations of motion for $Q_{f'}^\dagger$ and $\tilde{Q}^{\tilde{f}'\dagger}$ and become

$$\begin{aligned} \bar{D}^2(Q_{f'}^\dagger e^V Q^f) &= S \delta_{f'}^f + \hat{M}_{\tilde{f}}^f m_{f'}^{\tilde{f}} + b \hat{B} \delta_{f'}^f \\ \bar{D}^2(\tilde{Q}_{\tilde{f}} e^V \tilde{Q}^{\tilde{f}'\dagger}) &= S \delta_{\tilde{f}}^{\tilde{f}'} + m_{\tilde{f}}^{\tilde{f}'} \hat{M}_{\tilde{f}}^f + \tilde{b} \hat{\tilde{B}} \delta_{\tilde{f}}^{\tilde{f}'} \end{aligned} \quad (8)$$

respectively. Taking expectation values in a supersymmetric vacuum will remove the left-hand-sides of these equations (or equivalently working in the chiral ring of operators modded out by $\bar{D}$ exact operators). We finally arrive at

$$\begin{aligned} \langle \hat{M}_{\tilde{f}}^f \rangle m_{f'}^{\tilde{f}} &= (\langle S \rangle - b \langle \hat{B} \rangle) \delta_{f'}^f \\ m_{\tilde{f}}^{\tilde{f}'} \langle \hat{M}_{\tilde{f}}^f \rangle &= (\langle S \rangle - \tilde{b} \langle \hat{\tilde{B}} \rangle) \delta_{\tilde{f}}^{\tilde{f}'} \end{aligned} \quad (9)$$

We can obtain another set of equations similar to the above set by noting that there is another, gauge invariant, conserved current. Define the pair of operators

$$\begin{aligned} \hat{B}_{f_1}^{a_1} &:= \frac{1}{(N-1)!} \epsilon^{a_1 \dots a_N} \epsilon_{f_1 \dots f_N} Q_{a_2}^{f_2} \dots Q_{a_N}^{f_N} = \hat{B}(Q^{-1})_{f_1}^{a_1} \\ \hat{\tilde{B}}_{a_1}^{\tilde{f}_1} &:= \frac{1}{(N-1)!} \epsilon_{a_1 \dots a_N} \epsilon^{\tilde{f}_1 \dots \tilde{f}_N} \tilde{Q}_{\tilde{f}_2}^{a_2} \dots \tilde{Q}_{\tilde{f}_N}^{a_N} = \hat{\tilde{B}}(\tilde{Q}^{-1})_{a_1}^{\tilde{f}_1}. \end{aligned} \quad (10)$$

Under $SU(N_c)$ these transform exactly as $\tilde{Q}_{\tilde{f}}^a$ and Q_a^f respectively and therefore in the classical conservation laws above we may replace $\tilde{Q}_{\tilde{f}}^a$ by $\hat{B}_{\tilde{f}}^a$ and Q_a^f by $\hat{\tilde{B}}_a^{\tilde{f}}$ to obtain new conservation laws². Unlike the cases above however, these new conservation laws are not

²Such generalizations have also been considered in [14]

anomalous (as there is no propagator for $Q^\dagger$ and $\tilde{Q}$ or for $\tilde{Q}^\dagger$ and Q). Including the terms arising from the superpotential we obtain

$$\begin{aligned}\bar{D}^2(Q_f^\dagger e^V \hat{\tilde{B}}^{\tilde{f}}) &= m_f^{\tilde{f}} \hat{\tilde{B}} + b \hat{\tilde{B}} \hat{\tilde{B}} (\hat{M}^{-1})_f^{\tilde{f}} \\ \bar{D}^2(\hat{B}_f e^V \tilde{Q}^{\tilde{f}\dagger}) &= m_f^{\tilde{f}} \hat{B} + \tilde{b} \hat{\tilde{B}} \hat{\tilde{B}} (\hat{M}^{-1})_f^{\tilde{f}}\end{aligned}\tag{11}$$

which may equivalently be written as

$$\begin{aligned}\bar{D}^2(Q_f^\dagger e^V \hat{\tilde{B}}^{\tilde{f}}) &= m_f^{\tilde{f}} \hat{\tilde{B}} + b \det \hat{M} (\hat{M}^{-1})_f^{\tilde{f}} \\ \bar{D}^2(\hat{B}_f e^V \tilde{Q}^{\tilde{f}\dagger}) &= m_f^{\tilde{f}} \hat{B} + \tilde{b} \det \hat{M} (\hat{M}^{-1})_f^{\tilde{f}}\end{aligned}\tag{12}$$

where now the determinant is taken over the flavor indices. Taking expectation values as before removes the left-hand-side.

It was shown in [9] that an expectation value of a product of gauge invariant, chiral superfields is equal to the product of the individual expectation values. Applying this to the expectation values of (12) results in the equations

$$\begin{aligned}m_f^{\tilde{f}} \langle \hat{\tilde{B}} \rangle + b \det \langle \hat{M} \rangle (\langle \hat{M} \rangle^{-1})_f^{\tilde{f}} &= 0 \\ m_f^{\tilde{f}} \langle \hat{B} \rangle + \tilde{b} \det \langle \hat{M} \rangle (\langle \hat{M} \rangle^{-1})_f^{\tilde{f}} &= 0.\end{aligned}\tag{13}$$

We must be careful in obtaining this result. We could have taken the expectation values of these same equations in the form of (11). Applying the factorization property of [9] would have resulted (after some trivial manipulations) in the same equations as in (9) although without the S dependent term. Taken together these two sets of equations would have implied that S must vanish, which is not correct.

Our goal now is to use these equations in order to solve for the low energy effective action $W_{eff}(S, M, B, \tilde{B})$. To do this we note that³

$$\begin{aligned}\frac{\partial W_{eff}}{\partial M_f^{\tilde{f}}} &= -m_f^{\tilde{f}} \\ \frac{\partial W_{eff}}{\partial B} &= -b \\ \frac{\partial W_{eff}}{\partial \tilde{B}} &= -\tilde{b}.\end{aligned}\tag{14}$$

Therefore if we can solve the above equations for the Lagrange multiplier fields in terms of the variables appearing in the effective action, then we can integrate to find the effective action up to an S dependent term.

To begin we note that (9) implies that

$$b \hat{B} = \tilde{b} \hat{\tilde{B}}\tag{15}$$

³To simplify the notation we shall not write the expectation value signs $\langle \cdots \rangle$ from now on.

and therefore that

$$\hat{B} = \tilde{b}\xi, \quad \hat{\tilde{B}} = b\xi \quad (16)$$

where we have introduced a new variable ξ . We may therefore rewrite (9) as

$$\hat{M}_{\tilde{f}}^f = \left(S - \frac{\hat{B}\hat{\tilde{B}}}{\xi}\right)(m^{-1})_{\tilde{f}}^f. \quad (17)$$

We have at this stage reduced the problem to solving for ξ in terms of the meson and baryon vevs.

To find an equation for ξ we need only eliminate the Lagrange multipliers $m_{\tilde{f}}^{\tilde{f}}$ and b (or $\tilde{b}$) from (13) using the expressions in (17) and (16). This results in a linear equation for ξ with solution

$$\xi = -\frac{\det \hat{M} - \hat{B}\hat{\tilde{B}}}{S}. \quad (18)$$

Substituting for ξ in (16) and (17) we find

$$\begin{aligned} b &= -S \frac{\tilde{B}}{\det M - B\tilde{B}} \\ \tilde{b} &= -S \frac{B}{\det M - B\tilde{B}} \\ m_{\tilde{f}}^f &= S \frac{\det M}{\det M - B\tilde{B}} (M^{-1})_{\tilde{f}}^f. \end{aligned} \quad (19)$$

Finally we can insert these expressions back into the derivative equations (14) for W_{eff} and integrate. This results in

$$W_{eff} = -S \ln \left(\frac{\det M - B\tilde{B}}{\Lambda^{2N}} \right) + f(S) \quad (20)$$

where $f(S)$ remains to be fixed and we have put in the scale Λ in order to make the argument of the logarithm dimensionless. $f(S)$ is just the Veneziano-Yankielowicz contribution

$$W_{VY}(S) = (N_c - N_f)S \left(-\ln \frac{S}{\Lambda^3} + 1 \right) \quad (21)$$

which vanishes in this case as $N_c = N_f$. Therefore setting $f(S)$ to zero and minimizing W_{eff} with respect to S results in the constraint

$$\det M - B\tilde{B} = \Lambda^{2N} \quad (22)$$

and moreover a vanishing superpotential. This is in agreement with the results of [18].

3 $N_f = N_c + 1$

The methodology in the case $N_f = N_c + 1$ is identical to the previous section, although the details are slightly more involved. As before the superpotential that we consider consists of Lagrange multiplier terms that fix the vevs of the gauge invariant variables, i.e.,

$$W = m_{\tilde{f}}^f (\hat{M}_{\tilde{f}}^f - M_{\tilde{f}}^f) + b^f (\hat{B}_f - B_f) + \tilde{b}_{\tilde{f}} (\hat{\tilde{B}}^{\tilde{f}} - \tilde{B}^{\tilde{f}}) \quad (23)$$

where the meson operator was defined before, the baryon operators are defined as

$$\hat{B}_f := \frac{1}{N!} \epsilon_{ff_1 \dots f_N} \epsilon^{a_1 \dots a_N} Q_{a_1}^{f_1} \dots Q_{a_N}^{f_N} \quad (24)$$

$$\hat{\tilde{B}}^{\tilde{f}} := \frac{1}{N!} \epsilon^{\tilde{f} \tilde{f}_1 \dots \tilde{f}_N} \epsilon_{a_1 \dots a_N} \tilde{Q}_{\tilde{f}_1}^{a_1} \dots \tilde{Q}_{\tilde{f}_N}^{a_N} \quad (25)$$

and the fields $m_{\tilde{f}}^f$, b^f , and $\tilde{b}_{\tilde{f}}$ are Lagrange multpliers.

Proceeding exactly as before, we must derive the “loop” equations analogous to those found in (9) and (13). In the case of (9) the steps are virtually identical, so we shall forego the details and simply record the result. After taking expectation values we find

$$\begin{aligned} \langle \hat{M}_{\tilde{f}}^f \rangle m_{\tilde{f}}^f &= (\langle S \rangle - b^g \langle \hat{B}_g \rangle) \delta_{\tilde{f}}^f + b^f \langle \hat{B}_{\tilde{f}'} \rangle \\ m_{\tilde{f}}^{\tilde{f}'} \langle \hat{M}_{\tilde{f}}^f \rangle &= (\langle S \rangle - \tilde{b}_{\tilde{g}} \langle \hat{\tilde{B}}^{\tilde{g}} \rangle) \delta_{\tilde{f}}^{\tilde{f}'} + \tilde{b}_{\tilde{f}} \langle \hat{\tilde{B}}^{\tilde{f}'} \rangle. \end{aligned} \quad (26)$$

The second set of equations follows analogously to those found in (13). Namely we note that Q_a^f transforms under $SU(N_c)$ in the same way as the operator

$$\hat{\tilde{B}}_a^{\tilde{f} \tilde{f}'} := \frac{1}{N!} \epsilon^{\tilde{f} \tilde{f}' \tilde{f}_2 \dots \tilde{f}_N} \epsilon_{aa_2 \dots a_N} \tilde{Q}_{\tilde{f}_2}^{a_2} \dots \tilde{Q}_{\tilde{f}_N}^{a_N} \quad (27)$$

and therefore for vanishing superpotential we may define a new conserved current $(Q_{\tilde{f}'}^\dagger e^V \hat{\tilde{B}}^{\tilde{f} \tilde{f}'})$, which is not anomalous. In a completely analogous way we note that $\tilde{Q}_{\tilde{f}}^a$ transforms under color in the same way as

$$\hat{B}_{\tilde{f} f'}^a := \frac{1}{N!} \epsilon_{\tilde{f} f' f_2 \dots f_N} \epsilon^{aa_2 \dots a_N} Q_{a_2}^{f_2} \dots Q_{a_N}^{f_N} \quad (28)$$

and therefore for vanishing superpotential one has the conserved current $(\hat{B}_{\tilde{f} f'} e^V \tilde{Q}^{\tilde{f} \tilde{f} \dagger})$. Reinstating the superpotential (23) and taking expectation values results in the equations

$$\begin{aligned} (N-1)! (m_{\tilde{f}}^{\tilde{f}'} \langle \hat{\tilde{B}}^{\tilde{f}'} \rangle - m_{\tilde{f}}^{\tilde{f}} \langle \hat{\tilde{B}}^{\tilde{f}} \rangle) + b^g \epsilon_{\tilde{f}_1 \dots \tilde{f}_{N-1} f g} \epsilon^{\tilde{f}_1 \dots \tilde{f}_{N-1} \tilde{f} \tilde{f}'} \langle \hat{M}_{\tilde{f}_1}^{f_1} \rangle \dots \langle \hat{M}_{\tilde{f}_{N-1}}^{f_{N-1}} \rangle &= 0 \\ (N-1)! (m_{\tilde{f}}^{\tilde{f}} \langle \hat{B}_{\tilde{f}'} \rangle - m_{\tilde{f}}^{\tilde{f}'} \langle \hat{B}_{\tilde{f}} \rangle) + \tilde{b}_{\tilde{g}} \epsilon^{\tilde{f}_1 \dots \tilde{f}_{N-1} \tilde{f} \tilde{g}} \epsilon_{\tilde{f}_1 \dots \tilde{f}_{N-1} f f'} \langle \hat{M}_{\tilde{f}_1}^{f_1} \rangle \dots \langle \hat{M}_{\tilde{f}_{N-1}}^{f_{N-1}} \rangle &= 0 \end{aligned} \quad (29)$$

where as before we have used the factorization property of [9].

Now we can proceed to solve these equations. Recall that the goal is to construct the low energy effective action $W_{eff}(S, M, B_f, \tilde{B}^{\tilde{f}})$ where W_{eff} satisfies the partial differential equations

$$\begin{aligned}\frac{\partial W_{eff}}{\partial M_{\tilde{f}}^f} &= -m_{\tilde{f}}^f \\ \frac{\partial W_{eff}}{\partial B_f} &= -b^f \\ \frac{\partial W_{eff}}{\partial \tilde{B}^{\tilde{f}}} &= -\tilde{b}_{\tilde{f}}.\end{aligned}\tag{30}$$

We can use the equations (26) and (29) to solve for $m_{\tilde{f}}^f$, b^f and $\tilde{b}_{\tilde{f}}$ in terms of S , $M_{\tilde{f}}^f$, B_f and $\tilde{B}^{\tilde{f}}$, and then integrate to find W_{eff} up to an S dependent term.

To proceed let's first rewrite (26) in the form

$$\begin{aligned}\hat{M}_{\tilde{f}}^f &= (S - b^g \hat{B}_g)(m^{-1})_{\tilde{f}}^f + b^f \hat{B}_{f'}(m^{-1})_{\tilde{f}}^{f'} \\ \hat{M}_{\tilde{f}}^f &= (S - \tilde{b}_{\tilde{g}} \hat{\tilde{B}}^{\tilde{g}})(m^{-1})_{\tilde{f}}^f + \tilde{b}_{\tilde{f}}(m^{-1})_{\tilde{f}'}^f \hat{\tilde{B}}^{\tilde{f}'}\end{aligned}\tag{31}$$

where we have dropped the expectation value signs and will for the remainder of this section. Taking the trace of this equation results in the identity

$$b^f \hat{B}_f = \tilde{b}_{\tilde{f}} \hat{\tilde{B}}^{\tilde{f}}.\tag{32}$$

Inserting this back into (31) leads to

$$b^f \hat{B}_{f'}(m^{-1})_{\tilde{f}}^{f'} = \tilde{b}_{\tilde{f}} \hat{\tilde{B}}^{\tilde{f}'}(m^{-1})_{\tilde{f}'}^f.\tag{33}$$

This equation is enough to imply that

$$\begin{aligned}\hat{B}_f &= \tilde{b}_{\tilde{f}} m_{\tilde{f}}^f \xi \\ \hat{\tilde{B}}^{\tilde{f}} &= m_{\tilde{f}}^f b^f \xi\end{aligned}\tag{34}$$

where we have introduced the unknown ξ . One further result that we need from (31) can be obtained by contracting it with either $\hat{B}_f$ or $\hat{\tilde{B}}^{\tilde{f}}$. After some simplification and use of the relations (34), we find

$$\begin{aligned}(m^{-1})_{\tilde{f}}^f \hat{\tilde{B}}^{\tilde{f}} &= \frac{1}{S} \hat{M}_{\tilde{f}}^f \hat{\tilde{B}}^{\tilde{f}} \\ \hat{B}_f (m^{-1})_{\tilde{f}}^f &= \frac{1}{S} \hat{B}_f \hat{M}_{\tilde{f}}^f.\end{aligned}\tag{35}$$

Using (34) and (35) we may rewrite (31) as

$$m_{\tilde{f}}^{\tilde{f}} = (S - \frac{\hat{B}_g \hat{M}_g^g \hat{\tilde{B}}^{\tilde{g}}}{S\xi})(\hat{M}^{-1})_{\tilde{f}}^{\tilde{f}} - \frac{1}{S\xi} \hat{\tilde{B}}^{\tilde{f}} \hat{B}_{\tilde{f}}. \quad (36)$$

Combined with (34) we have managed to express the Lagrange multipliers in terms of the desired set of variables up to the unknown ξ .

To solve for ξ we must now bring in the equations (29). Eliminating $m_{\tilde{f}}^{\tilde{f}}$ and b^g or $\tilde{b}_{\tilde{g}}$ via the relations (36) and (34) respectively results after a bit of algebra in a linear equation for ξ with solution

$$\xi = -\frac{\det \hat{M} - \hat{B}_g \hat{M}_g^g \hat{\tilde{B}}^{\tilde{g}}}{S^2}. \quad (37)$$

This finally allows us to express the Lagrange multipliers in terms of the desired variables. We can now integrate the equations (replacing the operators $\hat{M}$ etc. by their vevs) for W_{eff} (30) to find

$$W_{eff} = -S \ln \left(\frac{\det M - B_g M_g^g \tilde{B}^{\tilde{g}}}{\Lambda^{2N}} \right) + f(S) \quad (38)$$

up to the S dependent function $f(S)$. $f(S)$ is given by the Veneziano-Yankielowicz term (21) for $N_f = N_c + 1$. Substituting and integrating out S we find for the low energy effective superpotential

$$W_{eff} = -\frac{1}{\Lambda^{2N-3}} (\det M - B_g M_g^g \tilde{B}^{\tilde{g}}) \quad (39)$$

in agreement with the results of [18].

4 $N_f > N_c + 1$

The results obtained thus far are not new, only the methods to obtain them are somewhat novel. The more interesting cases in which to apply these methods are when the number of flavors is even larger, i.e., when $N_f > N_c + 1$. In this case one can hope to address questions about confinement, chiral symmetry breaking, and Seiberg duality [19]. While the methods used above are straightforward to apply to the cases of $N_f > N_c + 1$, the equations that one obtains are more complicated and difficult to solve. We shall outline the construction in this section and leave its possible solution to future work.

As before we consider a superpotential which consists solely of Lagrange multiplier terms enforcing the meson and baryon vevs,

$$W = m_{\tilde{f}}^{\tilde{f}} (\hat{M}_{\tilde{f}}^{\tilde{f}} - M_{\tilde{f}}^{\tilde{f}}) + b^{f_1 \dots f_{\tilde{N}}} (\hat{B}_{f_1 \dots f_{\tilde{N}}} - B_{f_1 \dots f_{\tilde{N}}}) + \tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} (\hat{\tilde{B}}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} - \tilde{B}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}}) \quad (40)$$

where we have defined $\tilde{N} := N_f - N_c = N_f - N$ and the baryonic operators

$$\hat{B}_{f_1 \dots f_{\tilde{N}}} := \frac{1}{N!} \epsilon_{f_1 \dots f_{\tilde{N}} g_1 \dots g_N} \epsilon^{a_1 \dots a_N} Q_{a_1}^{g_1} \dots Q_{a_N}^{g_N} \quad (41)$$

$$\hat{\tilde{B}}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} := \frac{1}{N!} \epsilon^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}} \tilde{g}_1 \dots \tilde{g}_N} \epsilon_{a_1 \dots a_N} \tilde{Q}_{\tilde{g}_1}^{a_1} \dots \tilde{Q}_{\tilde{g}_N}^{a_N}. \quad (42)$$

The remaining fields in the superpotential, $m_f^{\tilde{f}}$, $b^{f_1 \dots f_{\tilde{N}}}$ and $\tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}}$ are the Lagrange multipliers.

As before we first construct the Konishi anomaly equations which are given by

$$\begin{aligned} \langle \hat{M}_{\tilde{f}}^f \rangle m_{\tilde{f}'}^{\tilde{f}} &= (\langle S \rangle - b^{f_1 \dots f_{\tilde{N}}} \langle \hat{B}_{f_1 \dots f_{\tilde{N}}} \rangle) \delta_{\tilde{f}'}^{\tilde{f}} + \tilde{N} b^{f_1 \dots f_{\tilde{N}-1} f'} \langle \hat{B}_{f_1 \dots f_{\tilde{N}-1} f} \rangle \\ m_{\tilde{f}}^{\tilde{f}'} \langle \hat{M}_{\tilde{f}}^f \rangle &= (\langle S \rangle - \tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} \langle \hat{\tilde{B}}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} \rangle) \delta_{\tilde{f}}^{\tilde{f}'} + \tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}-1} \tilde{f}'} \langle \hat{\tilde{B}}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}-1} \tilde{f}} \rangle. \end{aligned} \quad (43)$$

The other pair of equations needed to solve for the low energy effective action follow in direct analogy to before. We define the operators

$$\begin{aligned} \hat{\tilde{B}}_a^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}+1}} &:= \frac{1}{N!} \epsilon^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}+1} \tilde{g}_2 \dots \tilde{g}_N} \epsilon_{a a_2 \dots a_N} \tilde{Q}_{\tilde{g}_2}^{a_2} \dots \tilde{Q}_{\tilde{g}_N}^{a_N} \\ \hat{\tilde{B}}_{f_1 \dots f_{\tilde{N}+1}}^a &:= \frac{1}{N!} \epsilon_{f_1 \dots f_{\tilde{N}+1} g_2 \dots g_N} \epsilon^{a a_2 \dots a_N} Q_{a_2}^{g_2} \dots Q_{a_N}^{g_N}. \end{aligned} \quad (44)$$

We note that these transform under the $SU(N_c)$ symmetry exactly as Q_a^f and $\tilde{Q}_f^a$ respectively and therefore for vanishing superpotential one has the exact conservation laws

$$\begin{aligned} \bar{D}^2(Q_f^\dagger e^V \hat{\tilde{B}}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}+1}}) &= 0 \\ \bar{D}^2(\hat{\tilde{B}}_{f_1 \dots f_{\tilde{N}+1}} e^V \tilde{Q}^{\tilde{f} \dagger}) &= 0. \end{aligned} \quad (45)$$

Putting the superpotential (40) back in and taking expectation values results in the equations

$$\begin{aligned} (N-1)! m_f^{\tilde{f}} (\langle \hat{\tilde{B}}^{\tilde{g}_1 \dots \tilde{g}_{\tilde{N}}} \rangle \delta_{\tilde{f}}^{\tilde{g}} - \sum_{j=1}^{\tilde{N}} \langle \hat{\tilde{B}}^{\tilde{g}_1 \dots \tilde{g}_{j-1} \tilde{g}_{j+1} \dots \tilde{g}_{\tilde{N}}} \rangle \delta_{\tilde{f}}^{\tilde{g}_j}) \\ + b^{f_1 \dots f_{\tilde{N}}} \epsilon_{f_1 \dots f_{\tilde{N}} p_1 \dots p_{N-1} f} \epsilon^{\tilde{g}_1 \dots \tilde{g}_{\tilde{N}} \tilde{q}_1 \dots \tilde{q}_{N-1} \tilde{g}} \langle \hat{M}_{\tilde{q}_1}^{p_1} \rangle \dots \langle \hat{M}_{\tilde{q}_{N-1}}^{p_{N-1}} \rangle = 0 \\ (N-1)! m_f^{\tilde{f}} (\langle \hat{\tilde{B}}_{g_1 \dots g_{\tilde{N}}} \rangle \delta_g^{\tilde{f}} - \sum_{j=1}^{\tilde{N}} \langle \hat{\tilde{B}}_{g_1 \dots g_{j-1} g_{j+1} \dots g_{\tilde{N}}} \rangle \delta_{g_j}^{\tilde{f}}) \\ + \tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} \epsilon^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}} \tilde{p}_1 \dots \tilde{p}_{N-1} \tilde{f}} \epsilon_{g_1 \dots g_{\tilde{N}} q_1 \dots q_{N-1} g} \langle \hat{M}_{\tilde{p}_1}^{q_1} \rangle \dots \langle \hat{M}_{\tilde{p}_{N-1}}^{q_{N-1}} \rangle = 0 \end{aligned} \quad (46)$$

where we have used the factorization property of gauge invariant, chiral operators in writing these equations.

The procedure for solving these equations for the Lagrange multipliers in terms of the meson and baryon expectation values is in principle straightforward. We begin by

multiplying (46) by $(\tilde{N}+1)$ $\hat{M}$ operators and contracting with the free upper/lower indices respectively. This produces a $\det \hat{M}$ operator in the second term. Next we use (43) to eliminate the $m_f^{\tilde{f}}$ Lagrange multiplier in (46). After some algebra we arrive at

$$\begin{aligned}
\tilde{b}_{\tilde{g}_1 \dots \tilde{g}_{\tilde{N}}} &= -\frac{1}{\tilde{N}! \det \hat{M}} \left(\left(S + \left(\frac{N_f}{N} - 2 \right) (b \cdot \hat{B}) \right) (\hat{B} \hat{M})_{\tilde{g}_1 \dots \tilde{g}_{\tilde{N}}} \right. \\
&\quad \left. - \left(\frac{N_f}{N} - 1 \right) \sum_{j=1}^{\tilde{N}} (b \cdot \hat{B})_{f'}^{\tilde{f}} (\hat{M}^{-1})_{\tilde{f}}^{\tilde{f}'} \hat{M}_{\tilde{g}_j}^{\tilde{f}'} (\hat{B} \hat{M})_{\tilde{g}_1 \dots \tilde{g}_{j-1} \tilde{f} \tilde{g}_{j+1} \dots \tilde{g}_{\tilde{N}}} \right) \\
b^{g_1 \dots g_{\tilde{N}}} &= -\frac{1}{\tilde{N}! \det \hat{M}} \left(\left(S + \left(\frac{N_f}{N} - 2 \right) (\tilde{b} \cdot \hat{\tilde{B}}) \right) (\hat{M} \hat{\tilde{B}})^{g_1 \dots g_{\tilde{N}}} \right. \\
&\quad \left. - \left(\frac{N_f}{N} - 1 \right) \sum_{j=1}^{\tilde{N}} (\tilde{b} \cdot \hat{\tilde{B}})_{\tilde{f}}^{\tilde{f}'} (\hat{M}^{-1})_{\tilde{f}}^{\tilde{f}'} \hat{M}_{\tilde{f}'}^{g_j} (\hat{M} \hat{\tilde{B}})^{g_1 \dots g_{j-1} f g_{j+1} \dots g_{\tilde{N}}} \right) \quad (47)
\end{aligned}$$

where we have introduced some notation to simplify the expression,

$$\begin{aligned}
(\hat{B} \hat{M})_{\tilde{g}_1 \dots \tilde{g}_{\tilde{N}}} &:= \hat{B}_{g_1 \dots g_{\tilde{N}}} \hat{M}_{g_1}^{g_1} \dots \hat{M}_{g_{\tilde{N}}}^{g_{\tilde{N}}} \\
(b \cdot \hat{B})_{f'}^{\tilde{f}} &:= b^{g_1 \dots g_{\tilde{N}-1} f} \hat{B}_{g_1 \dots g_{\tilde{N}-1} f'} \\
(b \cdot \hat{B}) &:= b^{g_1 \dots g_{\tilde{N}}} \hat{B}_{g_1 \dots g_{\tilde{N}}} \quad (48)
\end{aligned}$$

with the other quantities defined similarly.

At this stage we can eliminate one Lagrange multiplier, $b^{g_1 \dots g_{\tilde{N}}}$ say, to find a linear equation for $\tilde{b}_{\tilde{g}_1 \dots \tilde{g}_{\tilde{N}}}$ in terms of the meson and baryon expectation values. Unlike the previous cases however the resulting equation is considerably more difficult to solve. The key difference is that the last terms on the right-hand-sides of the equations in (47) were of the same form as the first terms on the right-hand-sides. In that case solving the equations was trivial because they could be reduced in the end to solving for a single variable. In this case no such reduction (as far as we know) is possible making solving the equations more involved.

One interesting point to note however is that the solution for the Lagrange multipliers is necessarily linear in the gluino condensate S . This is easy to see by simply scaling all Lagrange multipliers in (47) by S and noting that S then factors out of the equation. The same holds true in the equations for $m_f^{\tilde{f}}$ in (43). As a consequence, after adding in the Veneziano-Yankielowicz contribution, the low energy effective superpotential will take the form

$$W_{eff} = (N_c - N_f) S \left(-\ln \frac{S}{\Lambda^3} + 1 \right) + S f(M, B, \tilde{B}) \quad (49)$$

where $f(M, B, \tilde{B})$ denotes the contribution coming from solving the equations above.

5 Seiberg dual theories

When the number of flavors satisfies $N_c + 2 \leq N_f \leq (3/2)N_c$, the moduli space of the gauge theory discussed in the previous section is known to be singular at the point

$M_{\tilde{f}}^f = B_{f_1 \dots f_{\tilde{N}}} = \tilde{B}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} = 0$. It was conjectured by Seiberg that the singularity is due to the presence of particles becoming massless at this point in moduli space. Seiberg conjectured a dual IR description of this theory which moreover is valid at the origin of moduli space. This theory consists of N_f chiral matter fields q_f and $\tilde{q}^{\tilde{f}}$, a set of chiral scalar fields $\hat{M}_{\tilde{f}}^f$, and superpotential $W_m = \hat{M}_{\tilde{f}}^f \tilde{q}^{\tilde{f}} q_f$ where the gauge group is $SU(N_f - N_c)$ and the gauge indices have been suppressed.

There are several non-trivial consistency checks that are discussed in detail in [18] that this dual description must pass in order to have a chance to describe the same IR physics as the theory discussed in the previous section. First note that the global symmetries are the same. Secondly we note that the massless degrees of freedom away from the point $\hat{M}_{\tilde{f}}^f = 0$ are the same. To see this let us first define the dual mesonic and baryonic operators

$$\begin{aligned} \hat{N}_{\tilde{f}}^{\tilde{f}} &:= \tilde{q}^{\tilde{f}} q_f \\ \hat{B}_{f_1 \dots f_{\tilde{N}}}^{(m)} &:= \epsilon_{a_1 \dots a_{\tilde{N}}} q_{f_1}^{a_1} \dots q_{f_{\tilde{N}}}^{a_{\tilde{N}}} \\ \hat{\tilde{B}}_{(\tilde{m})}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} &:= \epsilon^{a_1 \dots a_{\tilde{N}}} \tilde{q}_{a_1}^{\tilde{f}_1} \dots \tilde{q}_{a_{\tilde{N}}}^{\tilde{f}_{\tilde{N}}} \end{aligned} \quad (50)$$

where the (m) super or subscript denotes that this is the dual baryonic operator. Because of the superpotential the dual meson is massive away from the point $\hat{M}_{\tilde{f}}^f = 0$. The baryonic operators of the dual theories have identical transformation properties, identifying them then the dual theories in the end are described by the same set of low energy degrees of freedom. Finally one can check that the 't Hooft anomaly matching conditions are satisfied.

To go even farther one would like to compute the low energy effective superpotential on both sides of the duality and show that they are the same. While we will be unable to do this we nevertheless think that it will be useful to use the analysis described above to find a set of equations whose solution would provide the low energy effective superpotential. The analysis is essentially identical to above, the only new features here are that in this case there is already a superpotential present, aside from the Lagrange multiplier terms that we will add, and also that this time around we have an extra set of scalar chiral matter fields to deal with which leads to more equations.

The superpotential that we will take is

$$\begin{aligned} W_m &= \hat{M}_{\tilde{f}}^f \hat{N}_{\tilde{f}}^{\tilde{f}} + m_{\tilde{f}}^{\tilde{f}} (\hat{M}_{\tilde{f}}^f - M_{\tilde{f}}^f) + b^{f_1 \dots f_{\tilde{N}}} (\hat{B}_{f_1 \dots f_{\tilde{N}}}^{(m)} - B_{f_1 \dots f_{\tilde{N}}}^{(m)}) \\ &+ \tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} (\hat{\tilde{B}}_{(\tilde{m})}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}} - \tilde{B}_{(\tilde{m})}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}}) \end{aligned} \quad (51)$$

where as before the Lagrange multipliers $m_{\tilde{f}}^{\tilde{f}}$, $b^{f_1 \dots f_{\tilde{N}}}$ and $\tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}}$ enforce the vev constraints. This form is valid provided that the vev of the scalar fields $\hat{M}_{\tilde{f}}^f$ is non-zero, which is all that we shall concern ourselves with here. The analysis now proceeds as before. In complete analogy to the previous sections (essentially one just replaces Q 's

with q 's) we find the set of Konishi anomaly equations (dropping expectation value signs as before)

$$\begin{aligned}\hat{M}_{\tilde{f}}^f \hat{N}_{f'}^{\tilde{f}} &= S\delta_{f'}^f - \tilde{N} b^{f_1 \dots f_{\tilde{N}-1} f} \hat{B}_{f_1 \dots f_{\tilde{N}-1} f'}^{(m)} \\ \hat{M}_{\tilde{f}}^f \hat{N}_f^{\tilde{f}'} &= S\delta_{\tilde{f}}^{\tilde{f}'} - \tilde{N} \tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}-1} \tilde{f}} \hat{\tilde{B}}_{(m)}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}-1} \tilde{f}'}\end{aligned}\quad (52)$$

and the set of conservation equations

$$\begin{aligned}\hat{M}_{\tilde{f}}^f \hat{\tilde{B}}_{(m)}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}-1} \tilde{f}} &= -\tilde{N}! b^{f_1 \dots f_{\tilde{N}-1} f} \hat{N}_{f_1}^{\tilde{f}_1} \dots \hat{N}_{f_{\tilde{N}-1}}^{\tilde{f}_{\tilde{N}-1}} \\ \hat{M}_{\tilde{f}}^f \hat{B}_{f_1 \dots f_{\tilde{N}-1} f}^{(m)} &= -\tilde{N}! \tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}-1} \tilde{f}} \hat{N}_{f_1}^{\tilde{f}_1} \dots \hat{N}_{f_{\tilde{N}-1}}^{\tilde{f}_{\tilde{N}-1}}.\end{aligned}\quad (53)$$

Finally we can get another set of equations by recalling that $\hat{M}_{\tilde{f}}^f$ is a chiral, gauge invariant scalar field. Taking the expectation value of its equation of motion leads to

$$m_{\tilde{f}}^{\tilde{f}} + \hat{N}_{\tilde{f}}^{\tilde{f}} = 0. \quad (54)$$

The strategy before was to solve for the Lagrange multipliers as functions of the low energy degrees of freedom $\hat{M}_{\tilde{f}}^f$, $\hat{B}_{f_1 \dots f_{\tilde{N}}}^{(m)}$ and $\hat{\tilde{B}}_{(m)}^{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}}$. This leads to a set of equations in complete analogy to (30) which can in principle be integrated to find W_{eff} up to an S -dependent term. As an example, we can easily use the equations above to construct an equation for $\tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}}}$. Multiply the second equation of (53) by an $\hat{M}$ on every free index. Then use (54) and (52) to replace the $\hat{N}$ dependence in terms of the desired variables. We find in the end

$$\begin{aligned}0 &= \hat{M}_{\tilde{f}_1'}^{f_1} \dots \hat{M}_{\tilde{f}_{\tilde{N}}'}^{f_{\tilde{N}}} \hat{B}_{f_1 \dots f_{\tilde{N}}}^{(m)} \\ &+ \tilde{N}! \tilde{b}_{\tilde{f}_1 \dots \tilde{f}_{\tilde{N}-1} \tilde{f}_{\tilde{N}}'} (S\delta_{\tilde{f}_1'}^{\tilde{f}_1} - \tilde{N}(\tilde{b} \cdot \hat{\tilde{B}}_{(m)})_{\tilde{f}_1'}^{\tilde{f}_1}) \dots (S\delta_{\tilde{f}_{\tilde{N}-1}'}^{\tilde{f}_{\tilde{N}-1}} - \tilde{N}(\tilde{b} \cdot \hat{\tilde{B}}_{(m)})_{\tilde{f}_{\tilde{N}-1}'}^{\tilde{f}_{\tilde{N}-1}}),\end{aligned}\quad (55)$$

i.e., a nonlinear equation for $\tilde{b}$ in contrast to the linear equation that we obtained previously (47) for the IR dual theory.

It is interesting to note that some information on the S -dependence of W_{eff} can be obtained without solving for W_{eff} explicitly. If we scale the Lagrange multipliers b and $\tilde{b}$ as well as the mesons $\hat{M}_{\tilde{f}}^f$ by S , then all S dependence divides out of (55) as well as from (52). Consequently we know that W_{eff} must take the form, up to an S -dependent constant of integration, $W_{eff} = Sg(M/S, B, \tilde{B})$ for some function g . Putting in the Veneziano-Yankielowicz contribution we find

$$W_{eff} = -N_c S \left(-\ln \frac{S}{\Lambda^3} + 1 \right) + Sg(M/S, B, \tilde{B}). \quad (56)$$

6 Matrix model approach

In light of the conjecture of Dijkgraaf and Vafa, it is natural to ask if the results obtained earlier can be obtained from a matrix model. This question has been discussed by various authors [24, 21, 25, 26]. Recall that the Dijkgraaf-Vafa conjecture relates the planar diagrammatic contribution to the matrix model free energy to the low energy effective superpotential of the gauge theory. Moreover the matrix model action is given by the gauge theory superpotential expressed in terms of the elementary superfields. The question becomes somewhat more subtle when baryons are involved, for two reasons. The first is that normally one extracts planar diagrams from the matrix model by taking a large N limit, that is, the planar diagrammatic contribution dominates in this limit with all planar diagrams scaling with N in the same way. When baryons are included, the diagrams that one would like to call “planar”⁴ no longer all scale with N in the same way, making their extraction complicated. A second reason that baryons are more troublesome here is that the previous discussion assumed that the number of flavors N_f was also taken large, i.e. $N_f = N + k$ where k is fixed, but it has been shown that when fundamental matter is included, only planar diagrams with at most a single boundary are relevant to the low energy effective superpotential. Taking N_f large will not help to isolate the planar diagrams with different numbers of boundaries. Nevertheless, it has been shown in [21] that the generalized planar diagrams (in the gauge theory) are the relevant ones that give rise in the end to the low energy effective superpotential. If it is also true that the generalized planar diagrams from the associated matrix model are relevant to computing W_{eff} , then extracting them remains another question.

In this section we derive a set of matrix model loop equations for arbitrary N_c and $N_f \geq N_c$. These equations have identical form to those constructed in previous sections provided that we identify the glueball field S with g_m/N where g_m is the matrix model coupling constant, except that we don’t have the factorization property that followed in the gauge theory case from chirality. Normally on the matrix model side this property follows from the large N limit, namely in the large N limit the disconnected diagrams dominate the correlator giving rise to factorization. If one can make sense of the large N limit in this case, then this might be good enough to derive the factored form of the loop equations thereby giving identical results to the gauge theory analysis above. However the naive large N limit does not produce sensible results as the different terms in the loop equation scale with N differently.

Deriving the matrix model loop equations is a straightforward matter. As an example we shall consider the $N_c = N_f$ case. Our matrix model is given by the integral

$$e^{-N^2 \mathcal{F}} = \int [dQ] [d\tilde{Q}] e^{-(N/g_m)S} \quad (57)$$

with action S given by the superpotential (1) considered previously in the $N_f = N_c$ case. Here the matrix model variables Q_a^f and $\tilde{Q}_{\tilde{f}}^a$ are $N \times N$ where $N_c = N_f = N$. To derive

⁴Some generalization of the definition of planar is needed here due to the presence of the ϵ tensor in the definition of a baryon, see eg. [21].

the loop equations we consider first the identity

$$0 = \int [dQ] [d\tilde{Q}] \frac{\partial}{\partial Q_a^f} \left(Q_a^{f'} e^{-(N/g_m)S} \right). \quad (58)$$

Expanding this out results in the equation

$$\langle \hat{M}_{\tilde{f}}^f \rangle m_{f'}^{\tilde{f}} = (\langle S \rangle - b \langle \hat{B} \rangle) \delta_{f'}^{\tilde{f}} \quad (59)$$

where we have identified $S = g_m/N$. Replacing the explicit Q variables by $\tilde{Q}$ variables in (58) results in another loop equation of the same form in terms of the $\tilde{b}$ and $\hat{\tilde{B}}$ fields. These loop equations are in fact identical to those in the gauge theory case.

The second set of loop equations is obtained by considering the identity

$$0 = \int [dQ] [d\tilde{Q}] \frac{\partial}{\partial Q_a^f} \left(\frac{1}{N!} \epsilon_{aa_2 \dots a_N} \epsilon^{\tilde{f}\tilde{f}_2 \dots \tilde{f}_N} \tilde{Q}_{\tilde{f}_2}^{a_2} \dots \tilde{Q}_{\tilde{f}_N}^{a_N} e^{-(N/g_m)S} \right). \quad (60)$$

Expanding out the derivative results in

$$m_{\tilde{f}}^{\tilde{f}} \langle \hat{\tilde{B}} \rangle + b \langle \det \hat{M} (\hat{M}^{-1})_{\tilde{f}}^{\tilde{f}} \rangle = 0. \quad (61)$$

Exchanging the Q and $\tilde{Q}$ variables results in a similar type loop equation involving $\tilde{b}$ and $\hat{\tilde{B}}$. The important point to note here is that the second term is an expectation value of a product of operators. To reproduce the gauge theory results we must replace this with a product of expectation values of individual mesons.

As we said before, this factorization is usually accomplished by taking a large N limit, however the different terms in the above loop equations scale differently with N making this limit less than helpful. Perhaps there is a double scaling limit that one could take that preserves the form of the loop equations while giving rise to factorization.

Acknowledgements

We thank D.A. Lowe for comments and especially C-I. Tan for numerous discussions. This research is supported in part by DOE grant DE-FE0291ER40688-Task A.

References

- [1] K.A. Intriligator and N. Seiberg, Nucl. Phys. Proc. Suppl. 45BC (1996) 1, hep-th/9509066.
- [2] M.E. Peskin, (1997), hep-th/9702094.
- [3] N. Seiberg, Phys. Lett. B318 (1993) 469, hep-ph/9309335.
- [4] R. Dijkgraaf and C. Vafa, Nucl. Phys. B644 (2002) 3, hep-th/0206255.

- [5] R. Dijkgraaf and C. Vafa, Nucl. Phys. B644 (2002) 21, hep-th/0207106.
- [6] R. Dijkgraaf and C. Vafa, (2002), hep-th/0208048.
- [7] F. Ferrari, (2002), hep-th/0210135.
- [8] R. Dijkgraaf et al., (2002), hep-th/0211017.
- [9] F. Cachazo et al., JHEP 12 (2002) 071, hep-th/0211170.
- [10] N. Seiberg and E. Witten, Nucl. Phys. B426 (1994) 19, hep-th/9407087.
- [11] N. Seiberg and E. Witten, Nucl. Phys. B431 (1994) 484, hep-th/9408099.
- [12] N. Seiberg, (2002), hep-th/0212225.
- [13] B. Feng, (2002), hep-th/0212274.
- [14] A. Brandhuber et al., (2003), hep-th/0303001.
- [15] V. Balasubramanian et al., (2003), hep-th/0303065.
- [16] F. Cachazo, N. Seiberg and E. Witten, JHEP 04 (2003) 018, hep-th/0303207.
- [17] P. Kraus, A.V. Ryzhov and M. Shigemori, (2003), hep-th/0304138.
- [18] N. Seiberg, Phys. Rev. D49 (1994) 6857, hep-th/9402044.
- [19] N. Seiberg, Nucl. Phys. B435 (1995) 129, hep-th/9411149.
- [20] Y. Demasure and R.A. Janik, Phys. Lett. B553 (2003) 105, hep-th/0211082.
- [21] I. Bena, R. Roiban and R. Tatar, (2002), hep-th/0211271.
- [22] K. Konishi, Phys. Lett. B135 (1984) 439.
- [23] K.i. Konishi and K.i. Shizuya, Nuovo Cim. A90 (1985) 111.
- [24] R. Argurio et al., Phys. Lett. B553 (2003) 332, hep-th/0211249.
- [25] H. Suzuki, JHEP 03 (2003) 036, hep-th/0212121.
- [26] I. Bena et al., (2003), hep-th/0303115.